## Recombinant Human IL-23/IL-12B&IL-23A Protein

Catalog No.: RP01160 Recombinant

## **Sequence Information**

Background

Description

**Bio-Activity** 

**Basic Information** 

Species Gene ID Swiss Prot HEK293 cells 3593&51561P29460&Q9 NPF7

Tags No tag(IL12B)&N-his(IL23A)

#### Synonyms

IL-23 p19/IL-12 p40;IL23;SGRF ;IL-23 alpha & IL-12 beta

### **Product Information**

Source Purification HEK293 cells > 95% by SDS-PAGE.

#### Endotoxin

< 0.1 EU/µg of the protein by LAL method.

#### Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

#### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

#### Contact

€

www.abclonal.com

# **Storage** Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. <br/> > After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Active Recombinant Human IL-12B&IL-23A Protein is produced by HEK293 cells

Ser328&Arg20-Pro189) of human IL-23 (Accession #NP\_002178.2/NP\_057668.1)

expression system. The target protein is expressed with sequence (Ile23-

1.Measured by its binding ability in a functional ELISA.Immobilized Human IL-12B&IL-23A at  $5\mu$ /mL (100  $\mu$ L/well) can bind Human IL23R with a linear range

of 4-135 ng/ml.|2.Measured by its ability to induce IL-17 secretion by mouse

splenocytes. The ED<sub>50</sub> for this effect is 21.23-84.94 ng/mL,

1.17×10<sup>4</sup>~4.71×10<sup>4</sup> units/mg.


fused with a 6×His tag at the N-terminus(IL23A).

Avoid repeated freeze/thaw cycles.


corresponding to a specific activity of




## Validation Data



Recombinant Human IL-23/IL-12B&IL-23A Protein was resolved with SDS PAGE under reducing (R) and non-reducing (NR) conditions, showing single bands at 20(IL-23A),45-50(IL-12B) kDa and 65-70 kDa, respectively.



Immobilized Human IL-12B&IL-23A at 5 $\mu$ g/mL (100  $\mu$ L/well) can bind Human IL23R with a linear range of 4-135 ng/mL.



Recombinant Human IL-23/IL-12B&IL-23A induce IL-17 secretion by mouse splenocytes. The  $ED_{s0}$  for this effect is 21.23-84.94 ng/mL, corresponding to a specific activity of  $1.17 \times 10^4 \sim 4.71 \times 10^4$ units/mg.