



cDNA Library, *S. cerevisiae*, Log Phase

02-701 500 ng

This cDNA library (plasmid DNA) is constructed from *Saccharomyces cerevisiae*, strain S288C-derived poly(A)⁺ RNA at the log phase by the Linker-Primer method (Ref.1) by Prof. H. Nojima of Osaka University. This library is unidirectionally cloned by using the oligo (dT)₁₈ linker primer which contains the restriction enzyme site of *Not I*, and *Bam*HI (*Bgl*II)-*Sma* I adaptor.

The pLZ3 vector (shown below) used in this library can not replicate in *S. cerevisiae* but contains pUCori for replication in *E. coli*

Application

PCR screening of known or unknown gene: Prepare the primers for the known or unknown gene (cDNA) and amplify the gene by PCR from this library followed by cloning to an appropriate vector.

Standard amplifying conditions: 35 cycles of PCR reactions using 10-100 ng of cDNA as a template. (Change the quantity of template and the number of cycles depending on the expression rate of mRNA of the objective gene.)

Specification

Quantity: 500 ng (40 ng/ul, 13ul) in 10 mM Tris-HCl-1mM EDTA (pH 7.5)

Quality: 1) Number of independent clones: 3.6 x 10⁶

2) Average insert size : longer than 1 kb

Storage: -20°C

References: Construction of this library is described in Supplementary data of Ref.3


1. Kobori M *et al*"Large scale isolation of osteoclast-specific genes by an improved method involving the preparation of a subtracted cDNA library." *Genes Cells* 3: 459-475 (1998) PMID: [9753427](#)
2. Tanaka S and Nojima H "Nik1: a Nim1-like protein kinase of *S. cerevisiae* interacts with the Cdc28 complex and regulates cell cycle progression." *Genes Cells* 1, 905-921 (1996) PMID: [9077450](#)
3. Tougan T, Okuzaki D, Nojima H. Chum-RNA allows preparation of a high-quality cDNA library from a single-cell quantity of mRNA without PCR amplification. *Nucleic Acids Res.*, 36(15):e92, (2008) PMID:[18603591](#)

Note

- * This library is to be used only by the purchaser. It is not allowed to amplify and transfer the library to a third person.
- * Related products: human tissue specific cDNA libraries and cDNA libraries of model organisms (See [HP](#)).

to be continued...

Fig. Structure of pLZ3 and the restriction sites.



MCS(pLZ3)

CpoI(3) SauI(b) MluI(5)	AatII(3) BglII(5) AscI(5)	BalI(b)
PstI(3) SacI(3) ApaI(3) -----	-----	-----
SseI(3) -----	T7 Promoter	EcoRI(5) XbaI(5) AfI _{II} I(5) BstXI(5)

NNNCTGCA CCTGCAAGGAGCTCGAACGGGCCCTTAGGACCGTAAATACGACTCACTATAAGGGAATTGACGCTCTAGATCTTAAGGGCCCAAGGGGTTGGCCA
NNNG ACGTGGACCTCCTCGAGCTGGCCCGGAATCCCTGCCATTATGCTGAGTGATATCCCTTAAGCTGCAGATCTAGAATTCCGCGGTTCCCCAACCGGT

BstEII(5)	NheI(5)	-----	SwaI(3)	NruI(b)	SacII(3)
SnaBI(b) Dra _{II} I(3) ----- SceI(3)	-----	NotI(5) T3 promoter	-----	SphI(5) ----- PacI(3) ----- SacI(3)	

CCTGGTAACCAACGGGGTGGCTAGCTAGGGATAACAGGGTATATAGCGGCCGCCCTTAGTGAGGGTTAATTAAATCGTACGTGCGATTAATTAAACCGCGGTGGAGCT CAAT
GCACCATGGTQCCCACCGATCGATCCCTATTGTCCTTATATCGCCGGGGAAATCACTCCAAATTAAATTAGCATGCGCTAATTAAATTGGCGCCACC TCGACTTA

TCGCCCTATAGTGAGTCGTATTA -3'
AGCGGGATATCACTCAGCATAAT -5'